arXiv:1209.6621v1 [math.QA] 28 Sep 2012

Quantum McKay correspondence and global dimensions for fusion and module-categories associated with Lie groups

R. Coquereaux [1, 2]

September 2012

Abstract

Global dimensions for fusion categories defined by a pair (G, k), where G is a Lie group and k a positive integer, are expressed in terms of Lie quantum superfactorial functions. The global dimension is defined as the square sum of quantum dimensions of simple objects, for the category of integrable modules over an affine Lie algebra at some level. The same quantities can also be defined from the theory of quantum groups at roots of unity or from conformal field theory WZW models. Similar results are also presented for those associated module-categories that can be obtained via conformal embeddings (they are "quantum subgroups" of a particular kind). Some calculations use the correspondence existing between periodic quivers for simply-laced Lie groups and fusion rules for module-categories of type $SU(2)$.

Keywords: Lie groups; fusion categories; conformal field theories; quantum symmetries.

Classification: 81R50; 81T40; 18D10; 33E99.

1 Introduction

Let k be a positive integer and G a simple, compact Lie group and $Lie(G)$ its complex Lie algebra. We call $\mathcal{A}_k(G)$ the category of integrable modules of the Kac-Moody algebra associated with $Lie(G)$ at level k, see e.g. [11]. It is semi-simple, monoidal (use the fusion product), and modular. It is also equivalent [7, 14] to a category constructed in terms of representations of the quantum group G_q at root of unity $q = exp(\frac{i\pi}{N^\vee+k})$, where $N^\vee$ is the dual Coxeter number of G (take the quotient of the category of tilting modules by the additive subcategory generated by indecomposable modules of zero quantum dimension). These categories plays a key role in the Wess - Zumino - Witten models of conformal field theory.

One can associate a quantum dimension (or Perron-Frobenius dimension) μ to the objects of $\mathcal{A}_k(G)$, and since there are finitely many simple objects n, we can sum the square of their quantum dimensions $\mu(n)$ to define a global dimension $|\mathcal{A}_k(G)| = \sum_n \mu(n)^2$. One purpose of this paper is to obtain an explicit formula for this number, that generalizes the notion of order for a finite group. The formula obtained in the sequel involves a quantum version of the superfactorial where the product is done over q-integers associated with the exponents of the chosen Lie group. This function, that we call quantum Lie superfactorial of type G, is therefore a kind of generalization of the quantum Barnes function (for the latter, just take $G = SU(N)$).

Another purpose of this paper is to obtain an explicit formula for the global dimension of a large class of module-categories $\mathcal{E}$ associated with $\mathcal{A}_k(G)$, namely those modules that can be

---

[1] *IMPA, Instituto Nacional de Matemática Pura e Aplicada, Rio de Janeiro, Brasil.*

[2] *UMI CNRS-IMPA (2924), on leave from CPT, UMR 7332, Luminy, Marseille, France.*

obtained via conformal embedding of G, at level k in a larger Lie group J, at level 1 (see the text for details). Using another language, a module-category such as $\mathcal{E}$ is a "quantum subgroup" of the quantum group G_q, for q a root of unity. Such quantum subgroups come in two sorts, type 1 and type 2, and conformal embeddings give rise to (particular) quantum subgroups of type 1.

Some of the results described in the present article were presented at a conference held in Cordoba (Argentina), in 2010. The article [3], from the proceedings, gives proofs and examples but does not detail the calculations leading to our definitions for the quantum Lie superfactorial of type G, in particular when the latter is exceptional. Our proof of the formula for $|\mathcal{A}_k(G)|$ uses several known facts that will be recalled in the text but also an expression of the Weyl denominator (classical or quantum) that relies on explicit results for periodic quivers of roots associated with simple Lie groups G. The latter are here obtained in terms of structure coefficients (fusion numbers) for modules over the Grothendieck ring associated with the category $\mathcal{A}_k(SU(2))$. The quantum McKay correspondence relating ordinary ADE Dynkin diagrams and quantum subgroups of type $SU(2)$ has been known for many years and goes back to [2], but here we use it as a tool, in the sense that we exploit a correspondence previously described in [19] between periodic quivers for G (see for instance [16]) and an underlying quantum SU(2) theory. This little known relation is only briefly mentioned and used in [3], but no description of the corresponding methods is available, for this reason the discussion to be found in 6 can be of independent interest. This should make our paper reasonably self-contained.

Section 2 summarizes the main results concerning the global dimension of the fusion category $\mathcal{A}_k(G)$ and (a particular family of) associated modules categories $\mathcal{E}$. Sections 3 and 4 gives details and proofs, respectively for $\mathcal{A}_k(G)$ and for $\mathcal{E}$. The definition of the quantum Lie superfactorial, for all cases, including the non simply laced cases, is given at the end of section 3. Section 5 discusses a few examples and applications, as well as possible extensions of the present work. Section 6, devoted to the periodic quiver representation of the Weyl vector, uses the already mentioned correspondence between periodic quivers of roots, for simply-laced simple Lie groups, and module-categories of type $SU(2)$, but the discussion is extended to the non simply laced cases. This section is largely independent of the others and can be read independently; however, the corresponding results are needed at the end of section 3.

Remark: the global dimension of G at level k is formally equal to the square inverse of the Chern-Simon partition function[27], for connections on G-principal bundles on the sphere S^3. The topics discussed in this paper are therefore also related to differential geometry and to the theory of quantum invariants for knots.

2 Main results

If n is a real number and q is complex, there are two standard conventions for the definition of q-numbers ("quantum numbers"), leading to $[n]_q$ or to $[[n]]_q$. Both are recalled below.

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}} \quad \text{and} \quad [[n]]_q = \frac{1-q^n}{1-q}$$

In the present paper we mostly use the first convention, and sometimes omit the surrounding square bracket, setting $n_q = [n]_q$.

We shall need the following definitions:

Definition 1 *Let G be a simply-laced complex simple Lie group and let q be a complex number. Call $\mathcal{E}xp$ the multiset of exponents of G. We define the quantum Lie superfactorial of type G:*

$$\mathrm{sf}_G[q] = \prod_{s\in\mathcal{E}xp} [s]!_q$$

Here $[s]!_q$ denotes the quantum factorial of s, i.e., $[s]!_q = \prod_{n=1}^{n=s}[n]_q$.

For non-simply-laced cases (G not of type ADE), our definition of the quantum Lie superfactorial functions involves appropriate multiplicative factors that are given at the end of section 3.

For fusion categories of the kind studied in this paper, the global dimension is given by:

Theorem 1 *Let G a simple Lie group with rank r, Coxeter number N, dual Coxeter number $N^\vee$, and let k be a non-negative integer. The global dimension of the fusion category $\mathcal{A}_k(G)$ is given by*

$$|\mathcal{A}_k(G)| \doteq \sum_n \mu(n)^2 = \frac{(N^\vee+k)^r}{2^{rN}\,\Delta\left(\sin\dfrac{\pi}{N^\vee+k}\right)^{rN}\;(\mathrm{sf}_G[q])^2}$$

where $\mathrm{sf}_G[q]$ is the quantum Lie superfactorial of type G, Δ is the determinant of the fundamental quadratic form[1]*, and $q = \exp i\pi/(N^\vee+k)$.*

We then suppose that the fusion category $\mathcal{A}_k(G)$ is given, and we consider module-categories $\mathcal{E}_k(G)$ associated with it. As before, our purpose is to calculate their global dimensions. The simplest situation occurs when the module-category $\mathcal{E}_k(G)$ is associated with the conformal embedding of $Lie(G)$, at level k, into another Lie algebra $Lie(J)$, at level 1. The definition of conformal embeddings belongs to the lore of affine Lie algebras but we re-write it here in a way that uses only the properties of finite dimensional Lie algebras.

Definition 2 *Let G and J be simple Lie groups, and k, ℓ be positive integers. There is a conformal embedding of G at level k, in J at level ℓ, if the following three conditions are satisfied: 1) There is an embedding of the Lie algebra $Lie(G)$ into $Lie(J)$, 2) The Dynkin index of the embedding is equal to k, 3) If $N_G^\vee$ and $N_J^\vee$ are the dual Coxeter numbers of G and J, the following equality holds: $\frac{dim(G)\times k}{k+N_G^\vee} = \frac{dim(J)\times \ell}{\ell+N_J^\vee}$. We call c the common value of the last two expressions. When*[2] *the integer ℓ is not specified, it is understood that $\ell = 1$. If G is semi-simple, not simple, the same definition holds, but now $k = (k_i)$ is a multi-index, the quantity c_i is defined for each simple component G_i of G and the equality of central charges (condition 3) should hold for $c = \sum c_i$.*

Starting with such a conformal embedding $(Lie(G), k) \subset (Lie(J), \ell = 1)$, a known construction – that will be recalled in a latter section – allows one to determine a particular module-category $\mathcal{E}$ that, in some sense, "measures" the embedding.

Theorem 2 *Consider a conformal embedding of the simple or semi-simple Lie group G, at level k, in the simple Lie group J, at level 1. This embedding is associated with a module-category $\mathcal{E} = \mathcal{E}_k(G)$, with an action of the fusion category $\mathcal{A} = \mathcal{A}_k(G)$. The global dimension of $\mathcal{E}$ is given by:*

$$|\mathcal{E}| = \sqrt{|\mathcal{A}| \times |\mathcal{J}|}$$

where $|\mathcal{J}|$ is the global dimension of the fusion category $\mathcal{J} = \mathcal{A}_1(J)$.

3 Global dimension for the fusion category $\mathcal{A}_k(G)$

Although sometimes written differently, the results that we need and that are recalled in the next four subsections are standard. The hurried reader may directly jump to the fifth, and last, subsection, where the relation with (quantum) superfactorial functions is established.

---

[1] Calling L the Cartan matrix, $D = \mathrm{diag}(\delta_s)$ the diagonal symmetrizing matrix normalized in such a way that its largest entries are equal to 1 (equivalently, the long roots have norm 2), and $K = L^{-1}D$, then $\Delta = det(K)$ is the inverse of the index of the sublattice of long roots in the weight lattice. For simply laced cases, $\Delta = det(L)^{-1}$.

[2] This is what will be assumed from now on: we set $\ell = 1$.

The quantum Weyl formula and the quantum Weyl denominator

If a representation n exists at level k, its quantum dimension $\mu(n)$ at that level is given by the quantum version of the Weyl formula recalled below, with $q = exp(i\pi/(N+k))$. The classical dimension is recovered by taking $q = 1$. This expression is usually obtained from the theory of quantum groups or from the theory of affine Lie algebras, but we take it here as a definition. Supposing n irreducible, we use the same notation[3] for the representation and its highest weight.

$$\mu(n) = \prod_{\alpha>0} \frac{\langle n+\varrho, \alpha\rangle_q}{\langle \varrho, \alpha\rangle_q}$$

Here ϱ is the Weyl vector, α runs over the set of positive roots, and $\langle .., ..\rangle_q$ denotes the q-number $[\langle .., ..\rangle]_q$ associated with the inner product between the chosen weights (we do not write the surrounding square bracket). Using the $n_q = [n]_q$ convention to define the q-numbers $\langle \varrho, \alpha\rangle_q$, one has[4]:

Definition 3 *The quantum Weyl denominator of a Lie group G at level k is defined as $\mathcal{D}_W = \prod_{\alpha>0} \langle \varrho, \alpha\rangle_q$*

Action of $SL(2,\mathbb{Z})$ and the Kac-Peterson formulae

Action of the group $SL(2,\mathbb{Z})$ on the vector space spanned by the simple objects $m, n\ldots$ of the modular category $\mathcal{A}_k(G)$ is described by matrices S and T representing the two generators $\tau \mapsto -1/\tau$ and $\tau \mapsto \tau + 1$ of the modular group. This representation (actually a representation of the double cover), known by Hurwitz long ago [9] for $G = SU(2)$, is given, for arbitrary simple Lie groups, by the following formulae [12]:

$$S_{mn} = \frac{i^{\Sigma_+}\sqrt{\Delta}}{(N^\vee + k)^{r/2}} \sum_{w\in W} \epsilon_w \, e^{-\dfrac{2i\pi\langle w(m+\varrho), n+\varrho\rangle}{N^\vee + k}}, \; T_{mn} = e^{2i\pi\left[\dfrac{\langle m+\varrho, m+\varrho\rangle}{2(N^\vee+k)} - \dfrac{\langle \varrho, \varrho\rangle}{2N^\vee}\right]} \delta_{mn}$$

where $N^\vee$ is the dual Coxeter number, k is the level, w runs over the Weyl group W of G, ϵ_w is the signature, r is the rank of G, ϱ is the Weyl vector, Σ_+ is the number of positive roots (also equal to the sum of exponents, i.e., to $r\,N/2$ where N is the Coxeter number), and Δ is the determinant of the fundamental quadratic form. With these definitions one has $(S\,T)^3 = S^2 = \mathcal{C}$, the "charge matrix" satisfying $\mathcal{C}^2 = \mathbb{1}$. The matrix S is unitary and symmetric. The diagonal matrix T, which is related to the quadratic Casimir operator, will not be needed it in the sequel.

From the matrix element S_{00} to the quantum Weyl denominator

Let α be a positive root, ϱ the Weyl vector, and $q = exp(i\pi/(g+k))$. The q-scalar product $\langle \varrho, \alpha\rangle_q = (q^{\langle \varrho,\alpha\rangle} - q^{-\langle \varrho,\alpha\rangle})/(q - q^{-1})$, where $\langle ., .\rangle$ is the fundamental quadratic form, can be written

$$\langle \varrho, \alpha\rangle_q = (exp(\frac{\alpha}{2}) - exp(\frac{-\alpha}{2}))\,[\frac{2i\pi}{k+g}]\,/\,(2i\,\sin\frac{\pi}{g+k})$$

where e^λ is a formal exponential defined by $e^\lambda[\mu] = e^{\langle \lambda, \mu\rangle}$, for λ, μ, arbitrary elements of the space of weights. A standard manipulation (see for instance [5]) allows one to write

$$\prod_{\alpha>0} (e^{\alpha/2} - e^{-\alpha/2}) = \sum_{w\in W} \epsilon_w \, e^{w\varrho}$$

Taking $m = n = 0$, the trivial weight, in the Kac-Peterson formula for S_{mn}, and using the fact that the number of positive roots is $rN/2$, one obtains the following result (notice that the i factors cancel):

---

[3] On purpose, we use the same notation for integers (say n), so that, for $G = SU(2)$, an irreducible representation n of highest weight n has (classical) dimension $n+1$ and quantum dimension $\mu(n) = [n+1]_q$ at level k.

[4] The quantity $\mathcal{D}_W$ defined below would read differently if we were using another convention for q-numbers.

The element S_{00} of the modular matrix S and the quantum Weyl denominator $\mathcal{D}_W = \prod_{\alpha>0} \langle \varrho, \alpha \rangle_q$ are related as follows (although sometimes written differently, see for instance [5], this is certainly well known):

$$S_{00} = \frac{2^{\frac{rN}{2}} \sqrt{\Delta}}{(g+k)^{r/2}} \left(\sin \frac{\pi}{g+k} \right)^{\frac{rN}{2}} \mathcal{D}_W$$

Quantum dimensions, global dimension, and the S matrix

Straightforward manipulations on the quantum Weyl formula, analogous to the previous ones, lead to the following equality, which is sometimes used, in other contexts, to define quantum dimensions themselves:

$$\mu(n) = S_{n0}/S_{00}$$

Both numerator and denominator are positive real numbers, and unitarity of the S matrix implies $\sum_n |S_{n0}|^2 = 1$. In particular, the global dimension $|\mathcal{A}_k(G)|$, defined as sum of squares of the q-dimensions of its simple objects, can be expressed as:

$$|\mathcal{A}_k(G)| \doteq \sum_n \mu(n)^2 = \frac{1}{S_{00}^2}$$

From the expression for S_{00} obtained in the last subsection, one gets:

$$|\mathcal{A}_k(G)| = \frac{(g+k)^r}{2^{rN}\, \Delta \left(\sin \frac{\pi}{g+k} \right)^{rN} \mathcal{D}_W^2}$$

From the (quantum) Weyl denominator to (quantum) Lie superfactorials

The last step of the proof of *Theorem 1* finally relies on the following equality:

Lemma 1 *The quantum Weyl denominator of G is equal to the quantum superfactorial of type G. In other words : $\mathcal{D}_W = \mathrm{sf}_G[q]$.*
The functions $\mathrm{sf}_G[q]$ are defined at the end of this section.

Although a seemingly simple enough property, we could not find a precise reference, in the literature, besides [3], that would express, for all cases, the Weyl denominator of G in terms of superfactorial functions associated with exponents, like what we do in *Lemma 1*. However, in the case $G = A_r$ it is a familiar result that the Weyl denominator $\mathcal{D}_W$ is a product of successive factorials; this is clear from the Schur formula that can be used to calculate dimensions of irreducible representations described by Young tableaux. Writing the result in terms of exponents of the group $SU(r+1)$ is immediate since the latter are given by the consecutive integers $\{1, 2, \ldots r\}$. More generally, for classical Lie groups or Lie algebras (series A_r, B_r, C_r, D_r), the fact that one can express the (classical) Weyl denominator in terms of a product of factorials associated with the exponents seems to belong to the folklore, and the property is also hidden in the generalized hook dimension formula for representations of classical groups. Therefore, even if not phrased in the same way, we shall consider that the result for classical series is known, and we shall concentrate on the exceptional cases. We need explicit results both for the classical and for the quantum Weyl denominator.
We find convenient to gather the table of scalar products $\langle \varrho, \alpha \rangle$ (between the Weyl vector and the positive roots) in the periodic quiver of G, also called ribbon diagram of G [17], [20]. We shall obtain the results for the Weyl denominator by inspection, ie from its explicit determination in all cases. As announced in the introduction we shall use the quantum McKay correspondence in an operational way in the sense that we shall use the fusion numbers for module-categories of type $SU(2)$ to determine the periodic quiver representation of the (classical) Weyl vector,

for all Lie groups. Since these results, together with their relations with fusion numbers for module-categories of type $su(2)$, do not seem to be available elsewhere, and since the method used to obtain them may be of independent interest, we gather them in section 6.
What we need to complete our analysis is however a result for the *quantum* Weyl denominator, but this is easy since the latter is obtained from the classical one by replacing each scalar product $\langle \varrho, \alpha \rangle$ by the corresponding q-number $[\langle \varrho, \alpha \rangle]_q$. The expressions for quantum Lie superfactorials given below, and therefore also the final conclusion (*Theorem 1*), follow directly from the tables given in section 6.

Simply laced cases (ADE):

- $G = A_r \sim SU(r+1)$, $\mathcal{E} = \{1, 2, \ldots, r\}$, and $\mathrm{sf}_q(r) \doteq \mathrm{sf}_{A_r}[q] = \prod_{s=1}^{s=r} [s]!_q$.
- $G = D_r \sim SO(2r)$, $\mathcal{E} = \{1, 3, 5, \ldots N{-}3, N{-}1; N/2\}$, and[5] $\mathrm{sf}_{D_r}[q] = [N/2]!_q \prod_{s,odd=1}^{s=N-1} [s]!_q$.
- $G = E_6$ then $\mathrm{sf}_{E_6}[q] = [1]!_q\, [4]!_q\, [5]!_q\, [7]!_q\, [8]!_q\, [11]!_q$
- $G = E_7$ then $\mathrm{sf}_{E_7}[q] = [1]!_q\, [5]!_q\, [7]!_q\, [9]!_q\, [11]!_q\, [13]!_q\, [17]!_q$
- $G = E_8$ then $\mathrm{sf}_{E_8}[q] = [1]!_q\, [7]!_q\, [11]!_q\, [13]!_q\, [17]!_q\, [19]!_q\, [23]!_q\, [29]!_q$

Non simply laced cases:

- $G = B_r \sim SO(2r+1)$, then $\mathcal{E} = \{1, 3, 5, \ldots, 2r-1\}$ and $\mathrm{sf}_{B_r}[q] = \prod_{s \in \mathcal{E}} \widetilde{[s]!_q}$ where $\widetilde{[s]!_q} = [\frac{s}{2}]_q\, [s-1]_q\, [s-2]_q \ldots [3]_q\, [2]_q\, [1]_q$, and $\widetilde{[1]!_q} = [1/2]_q$
- $G = C_r \sim Sp(2r)$, then $\mathcal{E} = \{1, 3, 5, \ldots, 2r-1\}$ and $\mathrm{sf}_{C_r}[q] = \prod_{s \in \mathcal{E}} \widetilde{[s]!_q}$ where $\widetilde{[s]!_q} = \left[\frac{s}{2}\right]_q \left[\frac{s-1}{2}\right]_q \cdots \left[\frac{s-\frac{s-3}{2}}{2}\right]_q \left[s - \frac{s-1}{2}\right]_q \left[\frac{s-\frac{s+1}{2}}{2}\right]_q \cdots \left[\frac{2}{2}\right]_q \left[\frac{1}{2}\right]_q$ and $\widetilde{[1]!_q} = 1$.
- $G = F_4$, then $\mathcal{E} = \{1, 5, 7, 11\}$ and
  $\mathrm{sf}_{F_4}[q] = \left[\frac{1}{2}\right]_q^2 1_q^3 \left[\frac{3}{2}\right]_q [2]_q^3 \left[\frac{5}{2}\right]_q^2 [3]_q^3 \left[\frac{7}{2}\right]_q [4]_q^2 \left[\frac{9}{2}\right]_q [5]_q^2 \left[\frac{11}{2}\right]_q [6]_q [7]_q [8]_q$
- $G = G_2$, then $\mathcal{E} = \{1, 5\}$ and $\mathrm{sf}_{G_2}[q] = \left[\frac{5}{3}\right]_q \left[\frac{4}{3}\right]_q 3_q\, 2_q\, 1_q \left[\frac{1}{3}\right]_q$

Classical limits. The limit $\lim_{q \to 1} \mathrm{sf}_{A_r}[q]$ of the quantum Lie superfactorial of type A_r coincides with the superfactorial function $\mathrm{sf}(r)$ defined in [25]. More generally, taking the classical limit leads to the following integers or rationals[6]:

$$\mathrm{sf}(r) = \mathrm{sf}_{A_r} = \prod_{s=1}^{s=r} s! \qquad\qquad \mathrm{sf}_{B_r} = \frac{1}{2^r} \prod_{s,odd=1}^{2r-1} s!$$

$$\mathrm{sf}_{D_r} = (r-1)! \prod_{s,odd=1}^{2r-3} s! \qquad\qquad \mathrm{sf}_{C_r} = \frac{1}{2^{r(r-1)}} \prod_{s,odd=1}^{2r-1} s!$$

$$\mathrm{sf}_{E_6} = 2^{25} 3^{10} 5^5 7^3 11^1 \qquad\qquad \mathrm{sf}_{F_4} = \frac{1}{2^6 2^6}\, 1!\, 5!\, 7!\, 11! = 2^{15} 3^7 5^4 7^2 11^1$$

$$\mathrm{sf}_{E_7} = 2^{47} 3^{22} 5^{10} 7^6 11^3 13^2 17^1 \qquad\qquad \mathrm{sf}_{G_2} = \frac{1!\, 5!}{3^3} = 40/9$$

$$\mathrm{sf}_{E_8} = 2^{97} 3^{47} 5^{21} 7^{14} 11^8 13^6 17^4 19^3 23^2 29^1$$

Notice that for non–ADE cases, the scaling factor is $\prod_s (\delta_s)^{\gamma/2}$ where the coefficients δ_s are defined in footnote 1.

Relation with the Barnes G function. When the argument is an integer, the quantum Barnes G function is related to the quantum superfactorial as follows: $G_q(r+2) = \mathrm{Sf}_q(r)$, where $\mathrm{Sf}_q(r) = \prod_{s=1}^{s=r} [[s]]!_q$, with $[[s]]!_q = \prod_{n=1}^{n=s} [[n]]_q$, so that it differs from $\mathrm{sf}_q(r) = \prod_{s=1}^{s=r} [s]!_q$ by a pre-factor: $\mathrm{sf}_q(r) = q^{-(r+1)r(r-1)/6}\, G_{q^2}(r+2)$. The (classical, i.e., $q \mapsto 1$) Lie superfactorial of type $A_r \sim SU(r+1)$ coincides with the usual superfactorial.

---

[5] In the D_r case, $N = 2r - 2$, and when r is even, $N/2$ appears twice.

[6] These numbers and sequences have been added to the OEIS [24]

4 Global dimension for a class of module-categories $\mathcal{E}$

To say that we have a module-category $\mathcal{E}$ associated with a monoidal category $\mathcal{A}$ amounts to say [21] that we are given a monoidal functor from $\mathcal{A}$ to the category of endofunctors of an abelian category $\mathcal{E}$. In this setting, we have also a centralizer or endomorphism category $\mathcal{O}(E) = End_{\mathcal{A}}\mathcal{E}$ whose Grothendieck ring is often called "Ocneanu algebra". The latter is naturally a bimodule over the Grotendieck ring of $\mathcal{A}$ (the "fusion algebra"). Assuming, as before, that all our categories are semi-simple, one can encode this bimodule structure with a collection of matrices ("toric matrices") W_{xy} defined by $m\,x\,n = \sum_y (W_{xy})_{m,n}\, y$, where $m, n, \ldots$ are simple objects of $\mathcal{A}$ and $x, y, \ldots$ simple objects of $\mathcal{O}(E)$. In particular, to every module-category $\mathcal{E}$ is associated a matrix $Z = W_{00}$, where 0 is the unit object of $\mathcal{O}(E)$. The matrix Z is such that $Z_{00} = 1$, and if $\mathcal{A} = \mathcal{A}_k(G)$, which is a modular category, Z has the special property of being modular invariant, i.e., of commuting with the action of the modular group $SL(2, \mathbb{Z})$. This particular toric matrix Z (or its associated sesquilinear form) is often called in physics "the modular invariant", or "partition function". With the particular choice $\mathcal{E} = \mathcal{A}_k(G)$, the matrix Z is just the identity matrix. The other toric matrices W_{xy} can be physically interpreted in terms of twisted partition functions (presence of defects in boundary conformal field theories, see [22]). Their matrix elements are non-negative integers (this applies in particular to Z itself). In the sequel, we only need to know that to every module-category $\mathcal{E}$ over a modular category $\mathcal{A}_k(G)$ is associated a matrix Z that commutes with the action of $SL(2, \mathbb{Z})$.

We now assume that we are in the situation described in the discussion prior to *Definition 2*, so that we have a conformal embedding of G at level k in a Lie group J, at level 1. This determines a module-category $\mathcal{E}$ over the fusion category $\mathcal{A}_k(G)$ thanks to a construction that we do not need to recall because it will be enough, for our purpose, to show how the modular invariant matrix Z of $\mathcal{E}$ is obtained from J.

Notations. The group G and the level k being chosen, we simply call $\mathcal{A} = \mathcal{A}_k(G)$. We also set $\mathcal{J} = \mathcal{A}_1(J)$. We call $\widehat{G}$ the set of (classes of) simple objects n of $\mathcal{A}_k(G)$, $\widehat{J}$ the set of (classes of) simple objects $\hat{a}$ of $\mathcal{A}_1(J)$, and $\widehat{E}$ the set of (classes of) simple objects a of $\mathcal{E}$. At this point one should mention that $\widehat{G}$ and $\widehat{E}$ can be used as basis to explicitly describe the ring or module structures at hand in terms of matrices or graphs (the former being the adjacency matrices of the latter) called fusion graphs. For this reason, simple objects like $n, \hat{a}, a$ are sometimes called "vertices" of respective types G, J, E.

Trigonometric identities for $\mathcal{E}$

Call $Z = \sum_{m,n\in\widehat{G}} Z_{m,n}\; m \otimes n$ the partition function of $\mathcal{E}$. We shall need later the (known) identity:

$$\sum_{m,n\in\widehat{G}} \mu(m)\, Z_{m,n}\, \mu(n) = \sum_{n\in\widehat{G}} \mu(n)^2 \quad (= |\mathcal{A}|)$$

Indeed, from the expression of quantum dimensions of simple objects m, n of $\mathcal{A}$ in terms of matrix elements of the modular generator S, the left hand side reads $\sum_{m,n} S_{0m} Z_{mn} S_{0n}/S_{00}^2$. The matrix elements S_{0m} are real and S is symmetric, so that the l.h.s. reads $(SZS^\dagger)_{00}/S_{00}^2$, or $(ZSS^\dagger)_{00}/S_{00}^2$ since $SZ = ZS$ from modular invariance of Z. Unitarity of S together with the normalization condition $Z_{00} = 1$ imply that the l.h.s. is just $1/S_{00}^2$, hence the result.
The right-hand side of this equality being independent of Z (and therefore, of $\mathcal{E}$), we see that the previous identity gives rise to trigonometric-like identities for every choice of $\mathcal{E}$.

Induction-restriction functor and quantum dimensions

It is convenient to think of $\mathcal{A}/\mathcal{E}$ as a homogenous space, both discrete and quantum. Classically, the dimension of the space Γ_a of vector valued functions defined on the quotient of a finite group G by a subgroup E, and valued in a vector space a, can be calculated either trivially as $dim(a) \times |G/E|$ or non trivially, when a is a representation space for E, by decomposing this space of functions Γ_a (the space of sections of a vector bundle over G/E) into a sum of irreducible representations of G; this applies in particular to the case where a is the trivial representation $\underline{0}$ of E, so that we obtain a decomposition of the space $\mathcal{F}$ (actually the algebra) of complex valued functions over G/E, and $|\mathcal{F}| = dim(\Gamma_{\underline{0}})$. Here $\mathcal{F}$ refers both to a discrete set and to its algebra of functions.

In our situation, there is an induction-restriction functor [21], so that the above still makes sense. Simple objects a of $\mathcal{E}$ determine, by induction, objects Γ_a of $\mathcal{A}$ that are usually not simple and can be decomposed into simple objects n of $\mathcal{A}$: We write $\Gamma_a = \sum_{n \uparrow \Gamma_a} n$. Like in the classical situation, the (quantum) dimension of Γ_a can be calculated in two ways :

$$\mu(\Gamma_a) = \sum_{n \uparrow \Gamma_a} \mu(n) \quad \text{and} \quad \mu(\Gamma_a) = \mu(a) \times |\mathcal{A}/\mathcal{E}|$$

where $|\mathcal{A}/\mathcal{E}|$ is defined as $|\mathcal{A}|/|\mathcal{E}|$.

Let us mention the following results that will not be used in the sequel: there is a special object of $\mathcal{E}$, denoted $\underline{0}$, with quantum dimension equal to 1, and whose space of sections $\mathcal{F} \doteq \Gamma_{\underline{0}}$ plays the role of an algebra of functions over a "quantum space". $\mathcal{F}$ is an algebra (a Frobenius algebra [8]) in the category $\mathcal{A}$. When $a = \underline{0}$ the previous equation reads: $|\mathcal{F}| = |\mathcal{A}/\mathcal{E}|$. If $\mathcal{E}$ is, like here, associated with a conformal embedding, the set of its simple objects can be given a ring structure ("self-fusion") compatible with the $\mathcal{A}$ action, $\underline{0}$ playing the role of unity for this self-fusion. The modular invariant is then a sum of blocks (we are in the so-called "type I" case) and the decomposition of $\mathcal{F}$ into simple objects can be read from the first modular block of the partition function Z, or from the first line of the associated matrix.

A simple formula for $|\mathcal{E}|$

Because of the assumed conformal embedding of $Lie(G)$ into $Lie(J)$, objects of $\mathcal{J}$, and in particular simple objects (that we denote $\hat{a}$) can be considered as objects $\Gamma_{\hat{a}}$ of $\mathcal{A}$. The $\Gamma_{\hat{a}}$ are usually not simple and can be decomposed (using restriction, or "branching"), with multiplicities $\lambda_{\hat{a}}^n$, as a sum of simple objects n of $\mathcal{A}$. We write $\hat{a} \in \widehat{J} \subset ob(\mathcal{J}) \hookrightarrow \Gamma_{\hat{a}} \in obj(\mathcal{A})$, with $\Gamma_{\hat{a}} = \sum_n \lambda_{\hat{a}}^n \, n$. We shall only need two results from the theory of conformal embeddings; although maybe written differently, they belong to the lore of conformal field theory, and they have been known for quite a while:

- The partition function Z of the module-category $\mathcal{E}$, written as a sesquilinear form, is obtained from the partition function of $\mathcal{J}$ (the unit matrix) by merely replacing the simple objects $\hat{a}$ by their restriction $\Gamma_{\hat{a}}$ to $\mathcal{A}$.
- For every simple object $\hat{a}$ of $\mathcal{J}$, the object $\Gamma_{\hat{a}}$ of $\mathcal{A}$ obtained by restriction can also be obtained, by induction from some simple object (also called $\hat{a}$) of $\mathcal{E}$. This determines a subset – also denoted $\widehat{J}$ – the so-called "modular vertices" of the set of simple objects of $\mathcal{E}$. In other words, we can identify $\widehat{J}$ with a subset of $\widehat{E}$.

From the (trivial) modular invariant of $\mathcal{J}$, namely $\mathbb{1} = \sum_{\hat{a},\hat{b} \in \widehat{J}} \delta_{\hat{a},\hat{b}} \; \hat{a} \otimes \hat{b}$, we obtain $Z = \sum_{\hat{a},\hat{b} \in \widehat{J}} \delta_{\hat{a},\hat{b}} \; \Gamma_{\hat{a}} \otimes \Gamma_{\hat{b}}$ with $\Gamma_{\hat{a}} = \sum_{n \in \widehat{G}} \lambda_{\hat{a}}^n \, n$. The modular invariant Z is therefore a sum of blocks $Z^{\hat{a}}$ labelled by an indexing set $\widehat{J}$ that can be identified either with the set of simple objects of $\mathcal{J}$ or with a particular subset of the set $\widehat{E}$ of simple objects of $\mathcal{E}$. Going back to quantum dimensions, we see that the general trigonometric identity for Z recalled in 4 reads here:

$$|\mathcal{A}| = \sum_{\hat{a} \in \widehat{J}} (\mu(\Gamma_{\hat{a}}))^2$$

Using now the general formula relating $\mu(\Gamma_a)$ with $\mu(a)$, see section 4, one finds

$$|\mathcal{A}| = \sum_{\hat{a} \in \widehat{J}} \mu(a)^2 \times |\mathcal{A}/\mathcal{E}|^2$$

But $\sum_{\hat{a}\in\widehat{\mathcal{J}}} \mu(a)^2 = |\mathcal{J}|$ is the global dimension of $\mathcal{J}$, so that $|\mathcal{A}| = |\mathcal{J}| \times |\mathcal{A}|^2/|\mathcal{E}|^2$.

Hence the final result (*Theorem 2*): $|\mathcal{E}|^2 = |\mathcal{A}| \times |\mathcal{J}|$.

Since the right hand side refers to global dimensions of the fusion categories $\mathcal{A} = \mathcal{A}_k(G)$ and $\mathcal{J} = \mathcal{A}_1(J)$, it can be evaluated immediately thanks to the general formula of the previous section without having to compute the quantum dimensions of the simple objects of $\mathcal{E}$ (one does not even need to know how many there are).

In the case where we start from a conformal embedding of G into J, with J simple, but G only semi-simple, the above still makes sense. Writing $G \sim G_1 \times G_2$, there is a way to perform a reduction[7] with respect to one of the two factors (there are actually several possibilities at this point), and one ends up with a module-category $\mathcal{E}_1$ associated with, say, $|\mathcal{A}_{k_1}(G_1)|$. The partition function of $\mathcal{E}$ is a sum of blocks (type I situation), as we have seen, but but it may not be so for $\mathcal{E}_1$. The formula obtained for the global dimension of $\mathcal{E}$ can be applied, and it is an almost trivial matters to obtain the global dimension of $\mathcal{E}_1$ for the former.

5 Examples and applications

5.1 Examples for $\mathcal{A}_k(G)$ and applications

The main theorem giving $|\mathcal{A}_k(G)|$ leads, for every choice of G, to fully explicit formulae that can be expressed in terms of "elementary functions", since $[n]_q = \sin\left(\frac{\pi n}{k+N}\right)/\sin\left(\frac{\pi}{k+N}\right)$. For convenience, we remind the reader the values of Coxeter numbers N, dual Coxeter numbers $N^\vee$, and "long indices" Δ^{-1} for all Lie groups (the quantity Δ, used in the text, is the determinant of the fundamental quadratic form, i.e., the inverse of the long index).

	A_r	B_r	C_r	D_r	E_6	E_7	E_8	F_4	G_2
$N:$	$r+1$	$2r$	$2r$	$2r-2$	12	18	30	12	6
$N^\vee:$	$r+1$	$2r-1$	$r+1$	$2r-2$	12	18	30	9	4
$\Delta^{-1}:$	$r+1$	4	2^r	4	3	2	1	4	3

Using these values together with the definition of the quantum Lie superfactorial functions, one finds, for instance, that the global dimension of $A_{N-1} \sim SU(N)$ at level k reads:

$$|\mathcal{A}_k(SU(N))| = \frac{N(k+N)^{N-1}}{2^{N(N-1)} \prod_{s=1}^{N-1} \sin^{2(N-s)}\left(\frac{\pi s}{k+N}\right)}$$

whereas, in the case of G_2, it is:

$$|\mathcal{A}_k(G_2)| = \frac{3(k+4)^2}{2^{12} \sin^2\left(\frac{\pi}{3(k+4)}\right) \sin^2\left(\frac{\pi}{k+4}\right) \sin^2\left(\frac{4\pi}{3(k+4)}\right) \sin^2\left(\frac{5\pi}{3(k+4)}\right) \sin^2\left(\frac{2\pi}{k+4}\right) \sin^2\left(\frac{3\pi}{k+4}\right)}$$

These expressions can be used to study various kinds of limits. If we remember that $r + rN$ gives the dimension dim_G of the Lie group G, we obtain the following "classical limit" :

$$|\mathcal{A}_k(G)| \underset{k\to\infty}{\sim} \frac{k^{dim_G}}{(2\pi)^{rN}\,\Delta\,(\mathrm{sf}_G)^2}\ , \text{ in particular, } |\mathcal{A}_k(SU(N))| \underset{k\to\infty}{\sim} \frac{N \times k^{N^2-1}}{(2\pi)^{N(N-1)}\,\left(\mathrm{sf}(N-1)\right)^2}$$

The level-rank duality property for Lie groups of type $A_{N-1} \sim SU(N)$ was observed in [10]; for the coefficient S_{00} of the modular matrix S it implies: $\sqrt{N}\ S_{00}[A_{N-1}, k] = \sqrt{k}\ S_{00}[A_{k-1}, N]$. In terms of global dimensions we have therefore

$$\frac{|\mathcal{A}_k(SU(N))|}{|\mathcal{A}_N(SU(k))|} = \frac{N}{k}$$

---

[7] For example, when $G = SU(2)$, there is a module-category called E_7 associated with $\mathcal{A}_{18}(SU(2))$ that can be obtained from a semi-simple conformal embeddings $SU(2) \times SU(3) \subset E_8$ followed by a contraction (see for instance [4] for details).

This equality can be used to get simple enough expressions for global dimensions of $SU(N)$ at level k, when the rank $N-1$ is large and the level k reasonably small.
Notice that the above reads as a trigonometric identity for any choice a, b of positive integers:

$$2^{a(1-a)}(a+b)^{a-1}\prod_{s=1}^{b-1}[\sin(\frac{s\pi}{a+b})]^{2(b-s)} = 2^{b(1-b)}(a+b)^{b-1}\prod_{s=1}^{a-1}[\sin(\frac{s\pi}{a+b})]^{2(a-s)}$$

Using the previous asymptotic expression of $|\mathcal{A}_k(SU(N))|$ for large k, the duality relation for $SU(N)$ gives immediately another asymptotic expression when the level k is fixed and the rank $r = N-1$ goes to infinity:

$$|\mathcal{A}_k(A_r)| \underset{r\to\infty}{\sim} \frac{1}{(2\pi)^{k(k-1)}\,[\mathrm{sf}(N-1)]^2} \times r^{(k^2)}$$

Higher sums

The global dimension that was defined by summing squares of quantum dimensions $\mu(n)$ over the simple objects n of a category $\mathcal{A}_k(G)$ appears as a particular case, for $s=-2$, of a generalized Riemann Zeta function defined as $\zeta_G(s,k) = \sum_n \mu(n)^{-s}$. We did not address the problem of obtaining general formulae for this function when G is a simple Lie group and s is an arbitrary integer, but simple formulae should certainly exist. For instance, taking $G = SU(2)$, we know that $|\mathcal{A}_k(SU(2))| = \zeta_{SU(2)}(-2,k) = \frac{1}{2}(k+2)\csc^2\left(\frac{\pi}{k+2}\right)$, but we found some experimental evidence that for p integer, both even and positive, the quantity $\zeta_{SU(2)}(-p,k)$ is given, for levels $k > p/2-1$, by the expression $(k+2)\,4^{-p/2}\binom{p}{\frac{p}{2}}\sin^{-p}\left(\frac{\pi}{k+2}\right)$. When p is even but negative, more involved expressions can also be found (the overall numerical factor is no longer a simple binomial coefficient and the monomial $(k+2)$ that multiplies $\sin^{-p}\left(\frac{\pi}{k+2}\right)$ should be replaced by an appropriate polynomial of degree $-p$ in the variable k). We hope that the perspective of proving or generalizing these observations may trigger the interest of the reader.

5.2 Examples for $\mathcal{E}$ and applications

The list of conformal embeddings has been known for more than twenty years, see [1, 13, 23]. For the sake of illustration we shall only remind the reader the list of conformal embedding for $G = SU(g)$ Lie groups. There are three regular series and a few sporadic cases.

Regular series. The following conformal embeddings are respectively called antisymmetric, adjoint and symmetric (take $G = SU(N)$ in all cases):

k	$k=N-2$, $N\geq 4$	$k=N$, $N \geq 3$	$k=N+2$, $N\geq 2$
J	$SU(N(N-1)/2)$	$Spin(N^2-1)$	$SU(N(N+1)/2)$

Sporadic cases.

G	$SU(2)$		$SU(3)$		$SU(4)$	$SU(6)$	$SU(8)$		$SU(9)$
k	10	28	9	21	8	6	1	10	1
J	$Spin(5)$	G_2	E_6	E_7	$Spin(20)$	$Sp(10)$	E_7	$Spin(70)$	E_8

In order to apply the formula giving the global dimension of the category[8] $\mathcal{E}_k(G)$ associated to the chosen conformal embedding of G at level k into J (see theorem 2), one only needs to know $|\mathcal{A}_k(G)|$ and $|\mathcal{J}| = |\mathcal{A}_1(J)|$. Both being already given by the same general formula, there is nothing else to do. However, it may speed up calculations to determine the vector Q of quantum dimensions (with $\hat{J}$ components) and $|\mathcal{A}_1(J)|$ for all possible choices of J. This is left as an exercise to the reader, but the full answer can be found in [3]. Let us just mention that, for $J = SU(p)$, the irreducible representations that exist at level 1 are the fundamental ones ($p-1$ of them), and the trivial. All have quantum dimension 1, so that $Q = (1, 1, \ldots, 1)$, and $|\mathcal{A}_1(J)| = p$. The global dimensions of the first few $\mathcal{E}_k(SU(N))$ measuring regular conformal embeddings of type $SU(N)$ are then immediately obtained:

---

[8] sometimes called a "conformally exceptional quantum subgroup" of G

- Regular antisymmetric series ($k = N-2, N = 4, 5, \ldots$). Using $|\mathcal{J}| = N(N-1)/2$, one finds $|\mathcal{E}_{N-2}(SU(N))| = \left\{12, 20\left(2+\sqrt{2}\right), 60\left(5+2\sqrt{5}\right), 504\left(7+4\sqrt{3}\right), \ldots\right\}$
- Regular adjoint series ($k = N, N = 3, 4, 5, \ldots$). Using $|\mathcal{J}| = 4$, one finds $|\mathcal{E}_N(SU(N))| = \left\{12, 16\left(2+\sqrt{2}\right), 40\left(5+2\sqrt{5}\right), 288\left(7+4\sqrt{3}\right), \ldots\right\}$
- Regular symmetric series ($k = N+2, N = 2, 3, 4, 5, \ldots$). Using $|\mathcal{J}| = N(N+1)/2$, one finds $|\mathcal{E}_{N+2}(SU(N))| = \left\{6, 12\left(2+\sqrt{2}\right), 40\left(5+2\sqrt{5}\right), 360\left(7+4\sqrt{3}\right), \ldots\right\}$

Notice that for $G = SU(2)$ the only regular embedding is symmetric (it is at level 4, with $J = SU(3)$), the associated "conformally exceptional quantum subgroup" $\mathcal{E}_4(SU(2))$ is described by the fusion graph D_4 (the usual Dynkin diagram of $Spin(8)$), it has four simple objects with quantum dimensions $1, 1, 1, \sqrt{3}$, so that we could recover $|\mathcal{E}_4(SU(2))| = 6$ by summing squares. Still taking $G = SU(2)$ we see that we have also two sporadic embeddings, into $Spin(5)$ and G_2 : one recovers the exceptional $\mathcal{E}_{10}(SU(2))$ and $\mathcal{E}_{28}(SU(2))$ respectively described (and usually denoted!) by the fusion diagrams E_6 and E_8, i.e., the usual Dynkin diagrams. Using $|\mathcal{A}_1(Spin(5))| = 4$ and $|\mathcal{A}_1(G_2)| = \frac{1}{2}(5+\sqrt{5})$, together with $|\mathcal{A}_k(SU(2))| = (1/2)(k+2)/\sin^2(\pi/(k+2))$, one obtains the global dimensions $|E_6| = 4(3+\sqrt{3})$ and $|E_8| = \dfrac{1}{2}(15(3+\sqrt{5}) + \sqrt{30(65+29\sqrt{5})})$, that could also be directly obtained, in those relatively simple two cases, by summing six or eight squares.

Before ending this section, let us mention that the adjoint conformal embedding exists for all Lie groups (an embedding of G at level $k = N^\vee$, its dual Coxeter number, into $Spin(dim(G))$. For instance, the largest "conformally exceptional quantum subgroup" of the simple Lie group E_8 is obtained from the embedding of $\mathcal{A}_{30}(E_8)$ into $\mathcal{A}_1(Spin(248)) = \mathcal{A}_1(D_{124})$ whose global dimension is 4 (no calculation needed: the only representations of D_r that exist at level 1 are the trivial, the vectorial, and the two spinorial, all of q-dimension 1). The global dimension of the associated module-category is therefore: $|\mathcal{E}_{30}(E_8)| = 2\times\frac{60^4}{2^{120}\;[1]!_q[7]!_q[11]!_q[13]!_q[17]!_q[19]!_q[23]!_q[29]!_q\;\sin\left[\frac{\pi}{60}\right]^{120}}$ where one should use $q = \exp\frac{i\pi}{60}$ since $N = 30$ and $k = 30$.

6 The periodic quiver representation of the Weyl vector

The main result of the first section relies on an explicit determination of the Weyl vector, both in quantum and classical cases, or more precisely, on the collection of its scalar products with all roots of the chosen simple Lie group. Of course, this could be obtained by a variety of means, but it is more in the spirit of the present paper to obtain in this information by making use of the correspondence [2] existing between module-categories of $SU(2)$ type and ADE Dynkin diagrams. Using this correspondence, a formula giving the inner products of roots, for some chosen simply-laced simple Lie group, in terms of the fusion coefficients defining the action of the category $\mathcal{A}_k(SU(2))$ on an appropriate module-category $\mathcal{G}$, was given (using another language) in [19]. More generally, the weights of a simple Lie group G with Coxeter number $N = k+2$, thought of as integral functions on the space of roots, can be described in terms of such fusion coefficients. We introduce in the next subsections a set of matrices that precisely encode this information and are related both with the intertwiners of conformal field theory (those of a $SU(2)$ - WZW model) and with the so-called periodic Auslander-Reiten quiver. We shall study separately the simply and non-simply laced cases and finally give the resulting expressions for the Weyl vector, from which the stated results for the Weyl denominator follow. The quantum situation is handled without further ado by just replacing the value of each scalar product (for instance in the Weyl formula) by its q-number analog.

6.1 Periodic essential matrices (simply-laced cases)

The fusion category $\mathcal{A}_k(SU(2))$ possesses $k+1$ simple objects $(\underline{n}) \in \{(\underline{0}), (\underline{1}), (\underline{2}), \ldots, (\underline{k})\}$. The Grothendieck ring (the fusion ring) has one algebraic generator $\sigma = (\underline{1})$ and the unit is $(\underline{0})$. The other linear generators are obtained by the Chebyshev recurrence $(\underline{n}) = (\underline{n-1})\,\sigma - (\underline{n-2})$. Moreover, we have one relation: $(\underline{k+1}) = 0$ which implies $(\underline{k-1}) = (\underline{k})\,\sigma$. These result are well known and rely on the fact that the same recurrence (but not the extra relation $(\underline{k+1}) = 0$) holds for the representation theory of $SU(2)$ itself.

In the previous sections, the symbol $\mathcal{E}$ used to denote some module-category of type G (a Lie group), but in the present section we shall, on purpose, always denote the module-categories of type $SU(2)$ by the symbol $\mathcal{G}$. In other words, $\mathcal{G} = \mathcal{E}_{N-2}(SU(2))$. The chosen Dynkin diagram G is called the fusion diagram of $\mathcal{G}$ because its adjacency matrix (also called G) describes the action of the generator σ of the fusion ring of $SU(2)$ on the simple objects of $\mathcal{G}$: $\sigma \times a = \sum_b G_{a,b}\, b$. Indecomposable module-categories $\mathcal{G}$ of type $SU(2)$, i.e., associated with $\mathcal{A}_k(SU(2))$, are classified by simply laced Dynkin diagrams. This classification was first obtained by theoretical physicists in the framework of conformal field theories (CFT) (classification of modular invariant partition functions for the WZW models of type $SU(2)$, [2]). It was set in a categorical framework by [21, 15]. Because of this correspondence, to every vertex a of a simply laced[9] Dynkin diagram G with r vertices and Coxeter number N one can associate *both* a fundamental weight ω_a of a Lie group G, and a simple object a of an $SU(2)$ module-category $\mathcal{G}$ at level $k = N-2$ (this 2 stands for the dual Coxeter number of $A_1 \sim SU(2)$). When $\mathcal{G} = \mathcal{A}_k(SU(2))$ itself, the Dynkin diagram is A_{k+1}, and the choice of a vertex determines either a fundamental weight of $SU(k+2)$ or a simple object of $\mathcal{A}_k(SU(2))$ belonging to the ordered set $(\underline{0}), (\underline{1}), (\underline{2}), \ldots, (\underline{k})$. This vertex can also be referred to by its position (notice the shift by 1) in the list $(1, 2, \ldots, k+1)$.

The adjacency matrix of the chosen Dynkin diagram is symmetric since we assumed that it is simply laced (the Cartan matrix is $2 - G$). We then consider the $r \times r$ matrices $F_n = F_{(\underline{n-1})}$ defined by the following (Chebyshev) recurrence relation (that implies $F_2 = G$):

$$F_n = F_{n-1}.G - F_{n-2}, \qquad F_0 = 0, \qquad F_1 = \mathbb{1}$$

These matrices, sometimes called "nimreps" or "annular matrices" in CFT, encode the module-action of the fusion ring of $SU(2)$ at level k on the Grothendieck group of $\mathcal{G}$, which is automatically a $\mathbb{Z}_+$ module (its structure constants $(F_n)_{ab}$, labelled by simple objects a, b of $\mathcal{G}$, and defined by $(\underline{n})a = \sum_b (F_{(\underline{n})})_{ab}\; b$ are non negative integers). The matrices themselves are labelled by the simple objects of $\mathcal{A}_k(SU(2))$, i.e., by integers belonging to the prescribed range, but if we take instead $n \in \mathbb{N}$ we notice that the sequence F_n is periodic with period $2N$, where $N = k+2$ is the Coxeter number. The first $k+1$ members of the sequence have non-negative integer matrix elements, but $F_0 = F_N = F_{2N} = 0$, and the next $k+1$ members have non-positive matrix elements since they obey $F_{N+m} = -F_{N-m}$ for all positive integers m. From the family of square matrices F_n one defines the rectangular matrices $\mathfrak{r}_a = (\mathfrak{r}_a)_{nb} = (F_n)_{ab}$, where $a, b \in G$. If we take $1 \leq n \leq N-1$ we obtain r rectangular matrices $\mathfrak{r}_a$ (one for each vertex of the Dynkin diagram G) with $N-1$ lines and r columns called intertwiners in conformal field theory, or "essential matrices". If we allow the index n to run from 0 to $2N-1$ (or from 1 to $2N$), the obtained matrix elements are integers of both signs, but the obtained matrices should be considered as periodic in the vertical direction because of the the periodicity of matrices F_n. In our context, we shall take instead $0 \leq n \leq N-1$ (or $1 \leq n \leq N$), the essential matrices have now N lines, but the first line (or the last line) is null. More precisely, we proceed as follows.

ADE Dynkin diagrams are bipartite (they are trees), we declare the first vertex v of a long branch of the chosen diagram G to be even ($\partial\, v = 0$), and grade the other vertices accordingly. This defines a $\mathbb{Z}_2$ grading $a \in G \to \partial a \in \mathbb{Z}_2$ on the vertices. In particular, for $G = A_{N-1}$, we take $\partial(\underline{0}) = \partial 1 = 0$, $\partial(\underline{1}) = \partial 2 = 1$, and more generally $\partial(\underline{s}) = s\, mod\ 2$, so that the grading is also compatible with the module action. We now define, for any vertex a of G a rectangular matrix $\mathfrak{e}_a^+$, of size $r \times N$, and matrix elements $(\mathfrak{e}_a^+)_{nb}$, with $b \in G$ and $n \in \{1, 2, \ldots, N\}$:

---

[9] dual Coxeter number and Coxeter number coincide: $N^\vee = N$

$$\begin{array}{lrcl} \text{If} \quad \partial a = 0, & (\mathfrak{e}_a^+)_{nb} & = & (F_n)_{ab} \\ \text{If} \quad \partial a = 1, & (\mathfrak{e}_a^+)_{nb} & = & (F_{n-1})_{ab} \end{array}$$

The first line (resp. the last line) of $\mathfrak{e}_a^+$ is null when a is odd (resp. even). For an arbitrary weight $\omega = \sum_a \lambda_a \, \omega_a$, of G, with $\lambda_a \in \mathbb{Z}$ if ω is integral, we set $\mathfrak{e}^+(\omega) = \sum_a \lambda_a \, \mathfrak{e}^+(\omega_a)$, with $\mathfrak{e}^+(\omega_a) = \mathfrak{e}_a^+$. In particular, for the Weyl vector $\varrho = \sum_a \omega_a$, we obtain $\mathfrak{e}^+(\varrho) = \sum_a \mathfrak{e}_a^+$. Because of the horizontal $\mathbb{Z}_2$-grading (bipartition of the vertices of G) and vertical $\mathbb{Z}_2$-grading (Chebyshev recurrence formula for F_n) the matrix elements of $\mathfrak{e}_a^+$ vanish in position (n, b) when n and b are not both even, or both odd, for all possible choices of $a \in G$. For this reason, let us call $\mathcal{R}^+$ the discrete set obtained by keeping only those matrix elements (thought as functions: $a \in G \to (\mathfrak{e}_a^+)_{nb} \in \mathbb{Z}$) that are either simultaneously even or odd for both n and b. This discrete set with $r \times N/2$ elements can obviously be identified with the set of positive coroots[10] of the Lie group G since the cardinality of the set of all coroots is $r \times N$.

Using the action of some bipartite Coxeter element on the set of roots or coroots, a known construction, see [16], attaches to every simple Lie group a particular periodic quiver whose vertices are precisely associated with roots (or coroots). We do not need much from this construction here, but we shall use nevertheless the little known fact, both stated and generalized in [19], that the discrete set $\mathcal{R}^+$, constructed previously, provides a geometrical snapshot of (half of) this periodic quiver. This result can be precisely formulated as follows:

Let ω be an arbitrary weight associated with the simply laced Dynkin diagram G, of Coxeter number $N = k + 2$, then $\mathfrak{e}^+(\omega)$, with matrix elements $(\mathfrak{e}^+(\omega))_{nb}$ defined as above in terms of fusion coefficients of the category $\mathcal{A}_k(SU(2))$ acting on a module-category $\mathcal{G}$ also described by the graph G, gives the scalar products[11] between the weight ω and the positive coroots of G located at positions (n, b) of the table $\mathcal{R}^+$.

By allowing the range of indices n, in the previous definition of $\mathfrak{e}_a^+$, to be $\{1, 2, \ldots, 2N\}$ rather than $\{1, 2, \ldots, N\}$, we would obtain *periodic essential matrices* $\mathfrak{e}_a$ and recover all (positive and negative) roots $\mathcal{R}$, but in the present article we shall only need the positive ones.

Example E_6,

$$G = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix} \Longrightarrow \mathfrak{e}^+(\varrho) = \begin{pmatrix} 1 & & 1 & & 1 & \\ & 3 & & 3 & & 2 \\ 2 & & 7 & & 2 & \\ & 6 & & 6 & & 5 \\ 4 & & 10 & & 4 & \\ & 8 & & 8 & & 5 \\ 4 & & 11 & & 4 & \\ & 7 & & 7 & & 6 \\ 3 & & 9 & & 3 & \\ & 5 & & 5 & & 3 \\ 2 & & 4 & & 2 & \\ & 1 & & 1 & & 1 \end{pmatrix}$$



The Chebyshev recurrence relation using the adjacency matrix G of E_6 has a period $2N$, with $N = N^\vee = 12$, so that from the definition of matrices $\mathfrak{e}^+$, we obtain

$$\mathfrak{e}_1^+ = \begin{array}{cccccc} 1 & & 0 & & 0 & \\ & 1 & & 0 & & 0 \\ 0 & & 1 & & 0 & \\ & 0 & & 1 & & 1 \\ 0 & & 1 & & 1 & \\ & 1 & & 1 & & 0 \\ 1 & & 1 & & 0 & \\ & 1 & & 0 & & 1 \\ 0 & & 1 & & 0 & \\ & 0 & & 1 & & 0 \\ 0 & & 0 & & 1 & \\ & 0 & & 0 & & 0 \end{array}, \quad \mathfrak{e}_2^+ = \begin{array}{cccccc} 0 & & 0 & & 0 & \\ & 1 & & 0 & & 0 \\ 1 & & 1 & & 0 & \\ & 1 & & 1 & & 1 \\ 0 & & 2 & & 1 & \\ & 1 & & 2 & & 1 \\ 1 & & 2 & & 1 & \\ & 2 & & 1 & & 1 \\ 1 & & 2 & & 0 & \\ & 1 & & 1 & & 1 \\ 0 & & 1 & & 1 & \\ & 0 & & 1 & & 0 \end{array}, \quad \mathfrak{e}_3^+ = \begin{array}{cccccc} 0 & & 1 & & 0 & \\ & 1 & & 1 & & 1 \\ 1 & & 2 & & 1 & \\ & 2 & & 2 & & 1 \\ 1 & & 3 & & 1 & \\ & 2 & & 2 & & 2 \\ 1 & & 3 & & 1 & \\ & 2 & & 2 & & 1 \\ 1 & & 2 & & 1 & \\ & 1 & & 1 & & 1 \\ 0 & & 1 & & 0 & \\ & 0 & & 0 & & 0 \end{array}$$

---

[10] Since, in this section, G is simply laced, we could write "roots" instead

[11] As usual, the scalar product is relative to the euclidean structure determined by the inverse of the Cartan matrix

$$\mathfrak{e}_4^+ = \begin{array}{cccccc} 0 & & 0 & & 0 & \\ & 0 & & 1 & & 0 \\ 0 & & 1 & & 1 & \\ & 1 & & 1 & & 1 \\ 1 & & 2 & & 0 & \\ & 2 & & 1 & & 1 \\ 1 & & 2 & & 1 & \\ & 1 & & 2 & & 1 \\ 0 & & 2 & & 1 & \\ & 1 & & 1 & & 1 \\ 1 & & 1 & & 0 & \\ & 1 & & 0 & & 0 \end{array}, \quad \mathfrak{e}_5^+ = \begin{array}{cccccc} 0 & & 0 & & 1 & \\ & 0 & & 1 & & 0 \\ 0 & & 1 & & 0 & \\ & 1 & & 0 & & 1 \\ 1 & & 1 & & 0 & \\ & 1 & & 1 & & 0 \\ 0 & & 1 & & 1 & \\ & 0 & & 1 & & 1 \\ 0 & & 1 & & 0 & \\ & 1 & & 0 & & 0 \\ 1 & & 0 & & 0 & \\ & 0 & & 0 & & 0 \end{array}, \quad \mathfrak{e}_6^+ = \begin{array}{cccccc} 0 & & 0 & & 0 & \\ & 0 & & 0 & & 1 \\ 0 & & 1 & & 0 & \\ & 1 & & 1 & & 0 \\ 1 & & 1 & & 1 & \\ & 1 & & 1 & & 1 \\ 0 & & 2 & & 0 & \\ & 1 & & 1 & & 1 \\ 1 & & 1 & & 1 & \\ & 1 & & 1 & & 0 \\ 0 & & 1 & & 0 & \\ & 0 & & 0 & & 1 \end{array}$$

The scalar products between an arbitrary weight $\omega = \sum_a \lambda_a \omega_a$ and the simple positive coroots are encoded by the table $\mathfrak{e}^+(\omega) = \sum_a \lambda_a\, \mathfrak{e}_a^+$. In particular, for the Weyl vector itself, $\varrho = \sum_a \omega_a$, we obtain $\mathfrak{e}^+(\varrho) = \sum_a \mathfrak{e}_a^+$, hence the result. As discussed previously, we only list the relevant entries, i.e., those that are even for the total Z_2 grading. Finally, by multiplying all the values of $\mathfrak{e}^+(\varrho)$, one obtains $\mathcal{D} = 1!\,4!\,5!\,7!\,8!\,11! = \prod_{s \in \mathcal{E}xp(E_6)} s! = \mathrm{sf}_{E_6}$, as it was announced.

Remark: calling $\{\omega_a\}$ the basis of fundamental weights and $\{\beta_b\}$ the (dual) basis of simple coroots with respect to the scalar product $\langle \cdot, \cdot \rangle$, i.e., $\langle \omega_a, \beta_b \rangle = \delta_{ab}$, we have $\varrho = \sum_a \omega_a$, so that $\langle \varrho, \beta_a \rangle = 1$ for all $a \in G$. This determines the positions of simple coroots (among all positive coroots) on the vertices of the half-quiver $\mathcal{R}^+$. Arrows on a quiver, for instance on $\mathfrak{e}^+(\varrho)$, run from top to bottom and follow the pattern determined by the folded Dynkin diagram drawn on the side (it should be reflected every two steps when going downwards).

6.2 Periodic essential matrices (non simply-laced cases)

When the simple Lie group G or rank r is not simply-laced we still call G the non-symmetric matrix such that $2 - G$ is the Cartan matrix. G is the adjacency matrix of a graph that also characterizes the chosen group. We still define matrices $r \times r$ matrices F_n, with integer matrix elements, by the same Chebyshev recurrence relation as in the simply-laced case (section 6.1). The sequence is again periodic, of period $2N$, where N is the Coxeter number of G, which now differs from the dual Coxeter number $N^\vee$.

For Lie algebras G_2, F_4 and B_r we follow the convention of drawing the short roots to the right, with scaling factors[12] respectively equal to $(1, 1/3)$, $(1, 1, 1/2, 1/2)$, and $(1, 1, \ldots, 1, 1/2)$. For the Lie algebra C_r we draw short roots to the left, with scaling factors $(1/2, 1/2, \ldots, 1/2, 1)$. Remember that long roots have norm 2. Using the above drawing conventions, we attribute a Z_2 grading $\partial v = 0$ to the leftmost vertex v, and grade the other vertices accordingly, in an alternating way. We now can, in exactly the same way as in the simply-laced case, define matrices $\mathfrak{e}_a^+$, periodic essential matrices $\mathfrak{e}_a$, and more generally, matrices $\mathfrak{e}^+(\omega)$ or $\mathfrak{e}(\omega)$ for each weight ω of G. Again, keeping only the matrix elements of $\mathfrak{e}_a^+$ (thought as functions on fundamental weights) that are simultaneously even or odd for both gradings of lines and columns, one obtains a discrete set with $r \times N/2$ elements that can be identified with the set of positive roots, or with the set of positive coroots, of the group G. However, in the non simply-laced case, roots and coroots differ: the matrices $\mathfrak{e}_a^+$ just constructed give the scalar products between fundamental weights ω_a and all positive coroots (they are indeed non-negative integers) but the scalar products with the corresponding roots are obtained by introducing scaling factors for each orbit (columns of the quiver $\mathcal{R}^+$ i.e., vertical lines of matrices $\mathfrak{e}_a$), equal to the already recalled proportionality factors relating simple roots and simple coroots. These factors are non-trivial (not equal to 1) when the corresponding orbit is an orbit of short roots. We shall denote $\underline{\mathfrak{e}_a^+}$, and more generally $\underline{\mathfrak{e}^+(\omega)}$, for an arbitrary weight ω, the table encoding this information, i.e., the collection of scalar products between ω and all the positive roots. In particular, the Weyl denominator used in the first part of the paper is obtained by multiplying all the entries of $\underline{\mathfrak{e}^+(\varrho)}$.

Remark. For $SU(2)$ at level k, the duality $n \to \overline{n}$, corresponding to complex conjugation of representations, is trivial. As it is well known, this rules out the non-ADE Dynkin diagrams as describing possible fusion graphs for module-categories over $\mathcal{A}_k(SU(2))$ since the matrices describing the structure constants of the fusion ring of the latter, or its action on the associated modules, should be symmetric. Nevertheless, like in the ADE cases, the matrix elements of the F_n's (or, rather, of the $\mathfrak{e}_a$'s) still describe the scalar products between fundamental weights and coroots of a non simply-laced Lie group specified by the choice of the adjacency matrix G.

---

[12] The scalar product between fundamental weights is given by the elements of the (symmetric) matrix obtained by multiplying the inverse of the Cartan matrix with the diagonal matrix of scaling factors.

$$\textbf{Example } F_4\textbf{,} \quad G = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \to \mathfrak{e}^+(\varrho) = \left(\begin{array}{cccc} 1 & & 1 & \\ & 3 & & 2 \\ 2 & & 7 & \\ & 6 & & 5 \\ 4 & & 10 & \\ & 8 & & 5 \\ 4 & & 11 & \\ & 7 & & 6 \\ 3 & & 9 & \\ & 5 & & 3 \\ 2 & & 4 & \\ & 1 & & 1 \end{array}\right) \to \underline{\mathfrak{e}^+(\varrho)} = \left(\begin{array}{cccc} 1 & & 1/2 & \\ & 3 & & 2/2 \\ 2 & & 7/2 & \\ & 6 & & 5/2 \\ 4 & & 10/2 & \\ & 8 & & 5/2 \\ 4 & & 11/2 & \\ & 7 & & 6/2 \\ 3 & & 9/2 & \\ & 5 & & 3/2 \\ 2 & & 4/2 & \\ & 1 & & 1/2 \end{array}\right)$$



The Chebyshev recurrence relation using the adjacency matrix G of F_4 has a period $2N$, with $N = 12$ (here $N^\vee = 9$), so that from the definition of matrices $\mathfrak{e}^+$, we obtain

$$\mathfrak{e}_1^+ = \begin{array}{cccc} 1 & & 0 & \\ & 1 & & 0 \\ 0 & & 2 & \\ & 1 & & 2 \\ 1 & & 2 & \\ & 2 & & 0 \\ 1 & & 2 & \\ & 1 & & 2 \\ 0 & & 2 & \\ & 1 & & 0 \\ 1 & & 0 & \\ & 0 & & 0 \end{array}, \quad \mathfrak{e}_2^+ = \begin{array}{cccc} 0 & & 0 & \\ & 1 & & 0 \\ 1 & & 2 & \\ & 2 & & 2 \\ 1 & & 4 & \\ & 3 & & 2 \\ 2 & & 4 & \\ & 3 & & 2 \\ 1 & & 4 & \\ & 2 & & 2 \\ 1 & & 2 & \\ & 1 & & 0 \end{array}, \quad \mathfrak{e}_3^+ = \begin{array}{cccc} 1 & & 1 & \\ & 1 & & 1 \\ 1 & & 2 & \\ & 2 & & 1 \\ 1 & & 3 & \\ & 2 & & 2 \\ 1 & & 3 & \\ & 2 & & 1 \\ 1 & & 2 & \\ & 1 & & 1 \\ 0 & & 1 & \\ & 0 & & 0 \end{array}, \quad \mathfrak{e}_4^+ = \begin{array}{cccc} 0 & & 0 & \\ & 0 & & 1 \\ 0 & & 1 & \\ & 1 & & 0 \\ 1 & & 1 & \\ & 1 & & 1 \\ 0 & & 2 & \\ & 1 & & 1 \\ 1 & & 1 & \\ & 1 & & 0 \\ 0 & & 1 & \\ & 0 & & 1 \end{array}$$

The scalar products between the Weyl vector $\varrho = \sum_a \omega_a$ and the positive coroots are obtained as $\mathfrak{e}^+(\varrho) = \sum_a \mathfrak{e}_a^+$, and displayed above. However, the scalar products between the Weyl vector and the corresponding positive roots are obtained by scaling the last two columns of $\mathfrak{e}^+(\varrho)$ with a factor $1/2$. By multiplying all the values of $\underline{\mathfrak{e}^+(\varrho)}$, one obtains $\mathcal{D} = 1!\,5!\,7!\,11!/2^{12} = (1/2^{12}) \prod_{s \in \mathcal{E}xp(F_4)} s! = \mathrm{sf}_{F_4}$. Notice that the quantum Weyl denominator is obtained from the classical one by replacing the scalar products *themselves* by their q-analogs, so that for instance, $7/2$ becomes $[7/2]_q$, and this is not equal to $[7]_q/[2]_q$. By multiplying the q-analogs of all entries of the table $\underline{\mathfrak{e}^+(\varrho)}$, we recover the expression of the quantum superfactorial function of type F_4 that was given in section 3.

6.3 Other cases

Using the same method, and for every choice of the Lie group G, it is easy to determine the $\mathfrak{e}_a$, for all fundamental weights. Since we are only interested in the Weyl vector, and in the collection of its scalar products with positive roots, we only give the tables displaying $\mathfrak{e}^+(\varrho)$ when G is simply laced, and $\underline{\mathfrak{e}^+(\varrho)}$ when it is not. Results for the exceptional cases G_2, E_7 and E_8 are gathered in table 1 (the cases E_6 and F_4 have been described already). As explained in the last part of section 3, we focus our attention on the exceptional cases because, when G is not exceptional, the results needed to prove *lemma 1*, should be known. Nevertheless, we also display the cases A_6, B_6, C_6, D_6, in table 2 in order to illustrate what happens when G is a member of the classical series.

6.4 Miscellaneous remarks

General $\mathcal{G}$. The quiver interpretation using periodic essential matrices is not needed if our only goal is to determine the collection of scalar products entering the table $\mathfrak{e}^+(\varrho)$ since the same information is already contained in the matrices F_n. However, from their definition, one sees immediately that the coefficients of the table $\mathfrak{e}^+(\varrho)$ appear *twice* in the $r \times N$ matrix $(\sum_{b=1}^r ((F_n)_{a,b} + (F_{n-1})_{a,b}))$, indexed by (a, r) with $a = 1, \ldots r$ and $n = 1, \ldots N$. Therefore, the



Table 1: Values of the Weyl vector on the half-quiver of positive roots. Cases G_2, E_7, E_8.



A_6
1 2 2 1
3 4 3
1 5 5 1
3 6 3
1 4 4 1
2 2 2
D_6
1 2 3 2 2
3 5 5 4 1
1 6 7 7 3
4 8 9 6 1
1 3 5 4 2
1 3 5 4 2
B_6
1
2
1 2 2 2 2 2
3 4 4 4 4 1
1 5 6 6 6 3
3 7 8 8 5 1
1 5 9 10 7 3
$\frac{3}{2}$ $\frac{7}{2}$ $\frac{11}{2}$ $\frac{9}{2}$ $\frac{5}{2}$ $\frac{1}{2}$
C_6
2
1
$\frac{1}{2}$ 1 1 $\frac{3}{2}$ 1 1
$\frac{3}{2}$ 2 $\frac{5}{2}$ $\frac{5}{2}$ 2 $\frac{1}{2}$
$\frac{1}{2}$ $\frac{5}{2}$ $\frac{7}{2}$ $\frac{7}{2}$ $\frac{7}{2}$ $\frac{3}{2}$
$\frac{3}{2}$ 4 $\frac{9}{2}$ $\frac{9}{2}$ 3 $\frac{1}{2}$
$\frac{1}{2}$ 3 5 $\frac{11}{2}$ 4 2
2 4 6 5 3 1

Table 2: Values of the Weyl vector on the half-quiver of positive roots. Cases A_6, D_6, B_6, C_6.

square of the Weyl denominator of the simply laced simple Lie group with rank r and Coxeter number N defined by a Dynkin diagram G is obtained as:

$$\prod_{n=1}^{N} \prod_{a=1}^{r} \sum_{b=1}^{r} (F_n + F_{n-1})_{a,b} \;=\; \mathcal{D}^2 \;=\; (\mathrm{sf}_G)^2$$

where the F_n are the $r \times r$ matrices ("SU(2) nimreps") describing the module structure of the categorical module $\mathcal{G}$, specified by a fusion graph identical to the given Dynkin diagram, over the fusion algebra of $\mathcal{A}_k(SU(2))$ at level $k = N-2$, and where we also set $F_0 = F_N = 0$. The last term $n = N$ does not contribute to $\prod_{n=1}^{N}$, so that we could restrict the range to $\prod_{n=1}^{N-1}$. If the given simple Lie group is not simply laced, the same formula gives the square of the product, over all positive coroots, of the scalar product between ϱ and the positive coroots, so that in order to obtain a product over all positive roots one has to introduce known multiplicative factors (1/2 or 1/3) into the appropriate columns of the matrices $F_n + F_{n-1}$. The resulting overall multiplicative factor (with $q = 1$) appears in the definitions of the Lie superfactorial functions given at the end of section 3.

Special case $\mathcal{G} = \mathcal{A}_k(SU(2))$. In this particular case where $G = A_{k+1} \sim SU(k+2)$, the chosen module is the fusion algebra itself, the F matrices coincide with the fusion matrices and we may take all indices a, b and n from 1 to $r = k+1 = N-1$. The fusion coefficients $F_{p,m,n} = (F_p)_{m,n}$ are now completely symmetric with respect to permutation of their indices since the conjugation of representations is trivial for $SU(2)$. Using this property one can re-write the left hand side of the previous equation as $\prod_{n=1}^{r} \prod_{m=1}^{r} \sum_{p=1}^{r} (F_p + F_{p-1})_{m,n}$. Defining the path matrix[13] $X = \sum_{n=1}^{r} F_n$, and the *rotated* path matrix $\overset{\curvearrowleft}{X} = \lambda.X$, with λ the cyclic operator circularly permuting the lines, and using $F_0 = 0$, one obtains:

$$\prod_{m,n} (X + \overset{\curvearrowleft}{X})_{m,n} \;= 2^r \,(\mathrm{sf}(r))^2$$

With the previous definition of X, the above equality is valid for $\mathcal{G} = \mathcal{A}_k(SU(2))$ only.

Fourier-like considerations.

Relating the Weyl denominator of a simple Lie group, as we did, to matrices F_n describing modules over $SU(2)$ at level k, leads, if we diagonalize those matrices, to non-trivial identities relating trigonometric lines, sums of Chebyshev matrix polynomials, and superfactorial functions.

For a chosen module $\mathcal{G}$ over $\mathcal{A}_k(SU(2))$, equivalently for a simply laced Dynkin diagram G, the eigenvalues of the matrix $G = F_{(\underline{1})} = F_2$ are $\Delta_1 = 2\cos(\epsilon_j\, \pi/N)$, where ϵ_j, $\{j \in 1, \ldots, r\}$ belong to the multiset of exponents, denoted $\mathcal{E}xp$. Their set is included into the set of exponents of $\mathcal{A}_k(SU(2))$, ie into $\{1, \ldots, N-1\}$ with $k = N-2$. From this point of view the exponents of $\mathcal{G}$ (or of G) can be considered as a particular family of simple objects of $\mathcal{A}_k(SU(2))$, possibly coming with multiplicities. A more traditional approach to exponents defines them from the eigenvalues $\exp(2i\pi\epsilon_j/N)$ of some (arbitrary) Coxeter element, but the definition adopted here, using the adjacency matrix of a fusion graph (here a Dynkin diagram) and Perron-Frobenius theory, can be generalized to higher situations where $SU(2)$ is itself replaced by a more general Lie group.

Since the F_n matrices can be obtained from F_2 by a Chebyshev recurrence, they can be expressed as matrix polynomials $F_n = U_{n-1}(G/2)$, where $U_n(z) = \frac{1}{n!}\, \partial^n f[t,z]/\partial t^n \;|_{t=0}$ are the usual U-Chebyshev polynomials with generating function $f[t,z] = \frac{1}{t^2-2tz+1}$ that are such that $U_n[\cos\theta] = \sin((n+1)\theta)/\sin(\theta)$. For this reason, the F_n's constitute a commuting family of matrices with eigenvalues Δ_{n-1} with $\Delta_n = \sin((n+1)\epsilon_j\,\pi/N)/\sin(\epsilon_j\pi/N)$. To simplify the discussion, we assume, in the rest of this section, that the chosen Dynkin diagram G is simply laced (the matrix G is then symmetric). Calling Ψ an $r\times r$ unitary diagonalizing matrix for G, we write $F_n = \Psi^\dagger\,\mathrm{diag}(\Delta_n)\,\Psi$, where $\mathrm{diag}(\Delta_n)$ is a diagonal matrix with the above eigenvalues. Then

---

[13] X can be defined for an arbitrary $\mathcal{G}$ and the terminology comes from the fact that its elements give the dimensions of the vector spaces of essential paths [18],[28], on the fusion graph (also the dimensions of the vector spaces underlying the Gelfand-Ponomarev preprojective algebra associated with the corresponding unoriented quiver). The *sum* over columns of X gives the height vector. In the particular case $G = SU(N)$, X is a $r \times r$ matrix of increasing concentric rectangular rings of integers starting from 1 on the four sides.

$$\prod_{s=1}^{N}\prod_{a=1}^{r}\sum_{b=1}^{r}\left(\Psi^\dagger\ \mathrm{diag}\left[\frac{\sin\left(\frac{\pi\epsilon_j}{2N}+\frac{\pi s\epsilon_j}{N}\right)}{\sin\left(\frac{\pi\epsilon_j}{2N}\right)}\right]_{\epsilon_j\in\mathcal{E}xp}\Psi\right)_{a,b} = (\mathrm{sf}_G)^2$$

For instance, with $G = E_6$, $\mathcal{E}xp = \{1,4,5,7,8,11\}$ and Ψ as chosen below, the above expression is indeed equal to $(1!\,4!\,5!\,7!\,8!\,11!)^2$. For this example we set

$$\Psi = \frac{1}{2\sqrt{2}}\begin{pmatrix} \phi_- & \phi_+ & \sqrt{2}\,\phi_+ & \phi_+ & \phi_- & \sqrt{2}\,\phi_- \\ \sqrt{2} & \sqrt{2} & 0 & -\sqrt{2} & -\sqrt{2} & 0 \\ \phi_+ & \phi_- & -\sqrt{2}\,\phi_- & \phi_- & \phi_+ & -\sqrt{2}\,\phi_+ \\ \phi_+ & -\phi_- & -\sqrt{2}\,\phi_- & -\phi_- & \phi_+ & \sqrt{2}\,\phi_+ \\ \sqrt{2} & -\sqrt{2} & 0 & \sqrt{2} & -\sqrt{2} & 0 \\ \phi_- & -\phi_+ & \sqrt{2}\,\phi_+ & -\phi_+ & \phi_- & -\sqrt{2}\,\phi_- \end{pmatrix} \quad \text{with} \quad \phi_\pm = \sqrt{\frac{1}{3}\left(3\pm\sqrt{3}\right)}$$

In the particular case $\mathcal{G} = \mathcal{A}_k(SU(2))$, where the chosen module is the fusion algebra itself, one can take for Ψ the S-matrix itself representing one of the two generators of $SL(2,\mathbb{Z})$. Expressing the fusion matrices in terms of matrix elements of S alone is an identity known as the Verlinde formula [26]. Here we only need the matrix S for $SU(2)$, which is $S_{m,n} = \sqrt{\frac{2}{N}}\sin\left(\frac{\pi\,m\,n}{N}\right)$. Evaluation of the Weyl determinant then leads to the identity:

$$\prod_{s=1}^{N-1}\prod_{m=1}^{N-1}\sum_{n=1}^{N-1}\sum_{j=1}^{N-1}\frac{2}{N}\sin\left(\frac{\pi jm}{N}\right)\sin\left(\frac{\pi jn}{N}\right)\frac{\sin\left(\dfrac{\pi j}{2N}+\dfrac{\pi js}{N}\right)}{\sin\left(\dfrac{\pi j}{2N}\right)} = (\mathrm{sf}[N-1])^2 = \prod_{s=1}^{N-1}s!^2$$

References

[1] Bais F. and Bouwknegt P., A classification of subgroup truncations of the bosonic string, *Nucl. Phys.*, B 279, p 561, (1987).

[2] Cappelli A., Itzykson C. and Zuber J. -B., The ADE classification of minimal and $A_1^{(1)}$ conformal invariant theories, *Commun. Math. Phys.*, 13, pp 1–26, (1987).

[3] Coquereaux R., Global dimensions for Lie groups at level k and their conformally exceptional quantum subgroups, *Revista de la Union Matematica Argentina*, 51 No 2, pp 17-42 (2010).

[4] Coquereaux R., Conformal embeddings and quantum graphs with self-fusion. hal-00286087. Proceedings of the XVII Coloquio Latinoamericano de Algebra, Medellin, Colombia, July 2007.*S̃ao Paulo Journal of Mathematical Sciences* 3, 1 (2009), 239262

[5] Di Francesco P., Matthieu P. and Senechal D., Conformal field theory, Springer, (1997).

[6] Di Francesco P. and Zuber J.-B., SU(N) lattice integrable models associated with graphs, *Nucl. Phys.*, B 338, pp 602–646, (1990).

[7] Finkelberg, M., An equivalence of fusion categories, *Geom. Funct. Anal.* 6 (1996), 249 267.

[8] Fuchs J. and Schweigert C., Category theory for conformal boundary conditions, *Fields Institute Communications*, Vol. 39, pp 25-71, (2003).

[9] Hurwitz A,. Uber endliche Gruppen, welche in der Theorie der elliptschen Transzendenten auftreten Math. Annalen, 27, pp 183-233, (1886).

[10] Jimbo M. and Miwa T., A duality of branching rules for affine Lie algebras, *Adv. Stud. in Pure Math.*, 6, p 17, (1985).

[11] Kac V., Infinite dimensional Lie algebras, Cambridge University Press, Cambridge (1990).

[12] Kac V. and Peterson D., Infinite dimensional Lie algebras, theta functions, and modular forms, *Adv. Math.*, 53, pp 125–264, (1984).

[13] Kac V. and Wakimoto M. Modular and conformal invariance constraints in representation theory of affine algebras, *Adv. in Math.*, 70 p 156, (1988).

[14] Kazhdan D. and Lusztig G., Tensor structures arising from affine Lie algebras, III, *J. Amer. Math. Soc.*, 7, pp 335–381, (1994).

[15] Kirillov A. and Ostrik V., On q-analog of McKay correspondence and ADE classification of SL2 conformal field theories, *Adv. in Math.*, 171- 2, pp 183–227, (2002).

[16] Kirillov A. and Thind J., Coxeter elements and periodic Auslander – Reiten quiver, *Journal of Algebra*, 323 - 5, pp 1241-1265, (1 March 2010).

[17] Ocneanu A., seminars *unpublished*, (1998).

[18] Ocneanu A., Paths on Coxeter diagrams: from Platonic solids and singularities to minimal models and subfactors, Notes by Goto S., *Fields Institute Monographs*, Eds. Rajarama Bhat et al, (1999).

[19] Ocneanu A., The Classification of subgroups of quantum SU(N), in "Quantum symmetries in theoretical physics and mathematics", Bariloche 2000, Eds. Coquereaux R., García A. and Trinchero R., *AMS Contemporary Mathematics*, 294, pp 133–160, (2000).

[20] Ocneanu A., Higher Coxeter systems,
http://www.msri.org/publications/ln/msri/2000/subfactors/ocneanu, (2000).

[21] Ostrik V., Module categories, weak Hopf algebras and modular invariants, *Transform. groups*, 8, no 2, pp 177–206, (2003).

[22] Petkova V. B. and Zuber J.-B., The many faces of Ocneanu cells, *Nucl. Phys.*, B 603 , pp 449–496, (2001).

[23] Schellekens A. N. and Warner N. P., Conformal subalgebras of Kac-Moody algebras, *Phys. Rev.* D 34 - 10 , pp 3092 -3096, (1986).

[24] Sloane N. J. A., The On-Line Encyclopedia of Integer Sequences, `http://oeis.org`.

[25] Sloane N. J. A. and Plouffe S., The Encyclopedia of Integer Sequences, Academic Press, (1995).

[26] Verlinde E., Fusion rules and modular transformations in 2-D Conformal Field Theory, *Nucl. Phys.*, B 300, pp 360–376 (1988).

[27] Witten E., Quantum field theory and the Jones polynomial, *Comm. Math. Phys.*, 121, p 351, (1989).

[28] Zuber J.-B., CFT, BCFT, ADE and all that, Bariloche 2000, Eds. Coquereaux R., García A. and Trinchero R., *AMS Contemporary Mathematics*, 294, pp 133–160, (2000).